

# Tapping Gravitational Energy: Accretion onto Compact Stars

## Neutron Star Radii and Equation of State

# Contents

- Introduction
- Equation of State
- Comparing observations to theoretical EOS
- Neutron star cooling
- QPOs

# Properties

- Mass: 1-2 Solar Masses
- Density:  $\rho > 2\rho_0 = 2.7 \cdot 10^{14} \frac{\text{g}}{\text{cm}^3}$  in the core
- Rotation period: ms – hours, days
- Magnetic field:  $10^8 - 10^{14}$  G
- Radius:  $\sim 10$  km
- Gravitational acceleration:  $10^{13} - 10^{14} \frac{\text{cm}}{\text{s}^2}$

# Tolman-Oppenheimer-Volkov equation

$$\frac{dP(r)}{dr} = -\frac{G\mathcal{E}(r)M(r)}{c^2 r^2} \left(1 + \frac{P(r)}{\mathcal{E}(r)}\right) \left(1 + \frac{4\pi r^3 P(r)}{M(r)c^2}\right) \left(1 - \frac{2GM(r)}{c^2 r}\right)^{-1}$$

$$\frac{dM(r)}{dr} = 4\pi r^2 \rho(r) = \frac{4\pi r^3 \mathcal{E}(r)}{c^2}$$


$$M(r) = 4\pi \int_0^r R^2 \rho(R) dR = 4\pi \int_0^r R^2 \frac{\mathcal{E}(R)}{c^2} dR$$

# Equation of State

relationship between  $P(r)$  and  $\varepsilon(r) \Rightarrow P(\varepsilon)$

## 2 Assumptions:

- No rotation present, because rotation does not affect the fundamental EOS, it just changes the parameters we get up to about 20%
- Temperature is 0K. Reasonable because degeneration energy is much higher than the temperature




Pines 1991

# Atmosphere and Surface

- $\rho \leq 10^6 \frac{\text{g}}{\text{cm}^3}$
- Plasma of H, He, C, Fe in the atmosphere
- temperature and magnetic fields important for radiation
- atmosphere is not gray
- thin surface with large density gradient

# outer crust

- $10^6 \frac{\text{g}}{\text{cm}^3} \leq \rho \leq 4.3 \cdot 10^{11} \frac{\text{g}}{\text{cm}^3}$
- relativistic degenerate electron gas
- heavy, neutron rich nuclei
- $^{56}\text{Fe} \rightarrow ^{62}\text{Ni} \rightarrow ^{118}\text{Kr}$
- $\beta$ -equilibrium
- Coulomb lattice




# inner crust

➤  $4.3 \cdot 10^{11} \frac{\text{g}}{\text{cm}^3} \leq \rho \leq (2 - 2.4) \cdot 10^{14} \frac{\text{g}}{\text{cm}^3}$

➤ neutrons drop off from the nuclei

➤ superfluid neutrons




# neutron liquid

➤  $(2 - 2.4) \cdot 10^{14} \frac{\text{g}}{\text{cm}^3} \leq \rho \leq 2.8 \cdot 10^{14} \frac{\text{g}}{\text{cm}^3}$

➤ superfluid neutrons

➤ superconducting protons



# core region

➤  $2.8 \cdot 10^{14} \frac{\text{g}}{\text{cm}^3} \leq \rho$


➤ behavior of nuclear interactions not known

➤ muon condensation

➤ pion condensation

➤ hyperions

➤ quark matter



# Maximum mass (in general)


Assumptions:

- General relativity is valid
- microscopic stability  $\frac{dP}{d\rho} \geq 0$
- causality  $\frac{dP}{d\rho} \leq c^2$
- known EOS below some density  $\rho_0$


Result:

$$M_{\text{Max}} \approx 3.6 M_{\odot}$$

EOS	$M_{\max}$ [ $M_{\odot}$ ]	$R$ [km]	$r_g/R$	$n_c$ [fm $^{-3}$ ]	$\rho_c$ [ $10^{15}$ g cm $^{-3}$ ]
BPAL12	1.46	9.00	0.480	1.76	3.94
BGN1H1	1.64	9.38	0.519	1.60	3.72
BBB1	1.79	9.66	0.547	1.37	3.09
FPS	1.80	9.27	0.572	1.46	3.40
BGN2H1	1.82	9.53	0.564	1.45	3.48
BBB2	1.92	9.49	0.596	1.35	3.20
SLy	2.05	9.99	0.605	1.21	2.86
BGN1	2.18	10.9	0.592	1.05	2.26
APR	2.21	10.0	0.651	1.15	2.73
BGN2	2.48	11.7	0.626	0.86	2.02



P.Haensel, 2003




P.Haensel, 2003

# Cooling of Neutron Stars

- neutrino emission
  - dominates at least for the first 50000 years
  - Neutron star is transparent for neutrinos, because the their mean free path is much larger than the radius of a neutron star
- photon emission
  - photons are emitted only from the surface
  - radiation should have black-body spectrum

process	reaction	emissivity (erg/s/cm <sup>3</sup> )
direct URCA process	$n \rightarrow p + e^- + \bar{\nu}_e$ $p + e^- \rightarrow n + \nu_e$	$\sim 10^{27} \times T_9^6$
$\pi$ -condensate	$n + \pi^- \rightarrow n + e^- + \bar{\nu}_e$ $n + e^- \rightarrow n + \pi^- + \nu_e$	$\sim 10^{26} \times T_9^6$
quark URCA process	$d \rightarrow u + e^- + \bar{\nu}_e$ $u + e^- \rightarrow d + \nu_e$	$\sim 10^{26} \times T_9^6$
K-condensate	$n + K^- \rightarrow n + e^- + \bar{\nu}_e$ $n + e^- \rightarrow n + K^- + \nu_e$	$\sim 10^{24} \times T_9^6$
modified URCA process	$n + n \rightarrow n + p + e^- + \bar{\nu}_e$ $n + p + e^- \rightarrow n + n + \nu_e$	$\sim 10^{20} \times T_9^8$
coupled electron-neutrino processes	$\gamma + e^- \rightarrow e^- + \nu_e + \bar{\nu}_e$ $e^+ + e^- \rightarrow \nu_e + \bar{\nu}_e$	$\sim 10^{20} \times T_9^8$
Neutron-Neutron and Neutron-Proton Bremsstrahlung	$n + n \rightarrow n + n + \nu + \bar{\nu}$ $p + n \rightarrow p + n + \nu + \bar{\nu}$	$\sim 10^{19} \times T_9^8$
electron-ion-neutrino Bremsstrahlung	$e^- + (Z, A) \rightarrow e^- + (Z, A) + \nu_e + \bar{\nu}_e$	$\propto T_9^6$



- (1) RX J0822-4300
- (2) the Velar pulsar
- (3) PSR 0656+14
- (4) Geminga
- (5) RX J1856-3754
- (6) PSR 1055-52


- (a) Cas A point source
- (b) Crab pulsar
- (c) PSR J0205-6449
- (d) PSR 1509-58
- (e) PSR 1706-44
- (f) PSR 1823-13
- (g) PSR 2334+61
- (h) PSR 1951+32
- (i) PSR 0355+54
- (j) PSR 1929+10

- (A) 3C58

Tsuruta et al., 2002

# Remember QPO's


- two characteristic peaks
- assume sonic-point beat-frequency model
- frequency of the higher-frequency QPO is the Keplerian frequency at a special radius



Power spectrum of Sco X-1  
(Van der Klis, 2000)


## Getting maximum bounds for mass and radius of NS from QPOs:

- let 1220 Hz be the highest observed value of the higher-frequency
- the orbital radius  $R_{\text{orb}}$  of the clumps producing the QPO must be greater than the stellar radius
- $R_{\text{orb}}$  has also to be greater than the radius  $R_{\text{ms}}$  of the innermost stable circular orbit, in order to produce wave trains that lasts tens of oscillation periods



Lamp, Miller, Psaltis, 1998

Comparison the allowed mass/radius regions with some EOS




Lamp, Miller, Psaltis, 1998

## regions allowed for rotating NS and comparing with EOS

- consider a fixed frequency
- parameter for importance of rotational effects:

$$j \equiv \frac{cJ}{GM^2}$$

- for spin frequencies  $\sim 300\text{Hz}$   
j is about 0.1-0.3



Lamp, Miller, Psaltis, 1998

# Literature

- W. Becker, 1995, Ph.D. Thesis
- F.K. Lamb, M.C. Miller, D. Psaltis: Constraints on Neutron Star matter from kilohertz QPOs, 1998, astro-ph/9802348
- H. Heiselberg: Neutron Star Masses, Radii and Equation of State, 2002, astro-ph/0201465
- H. Heiselberg, M. Hjorth-Jensen: Phases of Dense matter in Neutron Stars, Physics Reports 328 (2000)
- P.Haensel: Equation of State of dense matter and maximum mass of Neutron Stars, 2003, astro-ph/0301073
- S. Tsuruta et al., Confronting Neutron Star Cooling Theories with New Observations, Astrophysical Journal Letters, Vol. 571, June 1, 2002
- F. Bocchino et al., The X-ray nebula of the filled center supernova remnant 3C58 and its interaction with the environment, 2001, astro-ph/0102087