



# Ali CMB Project

— An Introduction to **AliCPT**  
**(Ali CMB Polarization Telescope)**

**Xinmin Zhang**  
Institute of High Energy Physics  
Chinese Academy of Sciences  
On behalf of AliCPT collaboration

Dec. 18th, 2019 @ “B mode from space” , MPA, Munich

# Outline of the talk

- AliCPT site
- B mode physics (from space  $\oplus$  ground)
- AliCPT collaboration
- AliCPT current status, future plan and science
- Summary

# Where is the AliCPT (Ali CMB Polarization Telescope)

Located on a hilltop B1 ( $32^{\circ}18'38''$  N,  $80^{\circ}01'50''$  E),

in the Ngari (Ali) Prefecture of Tibet,

at an altitude of **5250 meters**.



- The Himalayas is to its southwest and runs from northwest to southeast, separating the Ngari(Ali) prefecture from the Indian Ocean.



- AliCPT (B1, 5250m) is only about **1km** to the current well-developed A1 station (5100m) of the Ali astronomical observatory.

# How to go to AliCPT



- Ngari Gunsa Airport is about 20 km away from the AliCPT site.
- Daily commercial flight between Ngari and Lhasa  
One flight between Ngari and Kashgar (Xinjiang Uyghur Autonomous Region) every Tuesday and Friday.



- The National Highway No. 219 from Airport to the Shiquanhe city;
- A road branched off to the AliCPT station at B1.



China National Highway G219

- Concrete pavement towards the B1 point has been completed.

Shiquanhe City



# Shiquanhe City

- Capital of Ali prefecture
- Population around 20,000,
- Equipped with culture, commerce and health facilities.



120 Emergency Center



Hotel (with Oxygen Provided)




Post office




It is only 30 km away from the AliCPT site.

# A1 station



Developed about 10 years ago  
150m lower than  
B1 (AliCPT)  
About 1km away from B1



- Some optical telescopes are running now.
- City grid **electric power** and the **network** connection have been set up.



# B1 station (AliCPT site)



**Altitude: 5250m**

**Coord : 32°18'38" N 80°01'50" E**

Figure left:

Construction began in Mar. 2017



Building 850m<sup>2</sup>

Weather station:


Pressure, wind speed/direction, temperature

City grid electric power / Generator

Network connection


Oxygen system

**Wind break**







# Ali: open up a new window for CMB polarization observations



Water vapor will:

- Reduce transmittance (absorption)
- Increase photon shot noise (emission)
- Increase optical loading on detectors (emission)






- AliCPT can cover
  - whole northern sky and partial southern sky
  - region with the lowest foreground contamination in the north.
- Overlap between AliCPT and southern experiments can be used for cross-check.
- Complementary to missions in Antarctic and Atacama. Realize full sky coverage.


South Pole, Atacama Desert, Ali

→ Full sky coverage


# CMB polarization: E/B mode




Scalar quadrupole,  
azimuthal symmetric  
generates T and E



Wayne hu



Tensor quadrupole,  
without azimuthal symmetry,  
generates T, E and B




- E mode: even Parity
- B mode: odd Parity

# The physics of the B mode

Observations:

CMB polarization maps




CMB-S4, arxiv: 1610.02743

$$\begin{array}{l} C_l^{BB} \\ C_l^{TB} \\ C_l^{EB} \end{array}$$

Estimate

Constrain



$$\begin{array}{l} r \\ \alpha \equiv \bar{\alpha} + \delta\alpha \end{array}$$


Phenomenological parameters:

- $r$ : tensor to scalar ratio
- $\bar{\alpha}$ : global CMB rotation angle
- $\delta\alpha$ : spatial dependent rotation angle

# Dynamical models for $r, \bar{\alpha}, \delta\alpha$

- $r$ : Inflation theory  
introducing  $\phi(x)$ , then inflaton

$$\mathcal{L} \sim \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi)$$




- $\alpha$  : Chern-Simons theory

$$V(\phi) \sim m^2 \phi^2, \lambda \phi^4, \dots$$

$$\mathcal{L}_{QED} \sim -\frac{1}{4} \mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu} + \frac{c}{M} \phi(x) \mathcal{F}_{\mu\nu} \tilde{\mathcal{F}}^{\mu\nu}.$$

where:  $\tilde{\mathcal{F}}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} \mathcal{F}_{\alpha\beta}$

$\phi(x)$ : Axion scalar,  
Axion string,  
Quintessence scalar



# Chern-Simons theory (Axion coupling)

$$\phi(x) \mathcal{F}_{\mu\nu} \tilde{\mathcal{F}}^{\mu\nu} \sim \phi(x) \partial_\mu K^\mu \longrightarrow \partial_\mu(\phi(x) K^\mu) - \partial_\mu \phi K^\mu$$


$$K^\mu \sim \epsilon^{\mu\nu\alpha\beta} A_\nu \mathcal{F}_{\alpha\beta}, \quad \int \partial_\mu(\phi(x) K^\mu) = 0 \quad (\text{Trivial vacuum})$$



$$\phi(x) \mathcal{F}_{\mu\nu} \tilde{\mathcal{F}}^{\mu\nu} \longrightarrow -\partial_\mu \phi(x) K^\mu$$

# During $\phi(x)$ evolution

$$\partial_\mu \phi(x) \longrightarrow \langle \partial_0 \phi(x) \rangle$$



**For RW universe, time dependent vacuum expectation.**

$$\partial_\mu \phi(x) K^\mu \longrightarrow \langle \partial_0 \phi(x) \rangle K^0$$

**Under CPT transformation:**

$$\langle \partial_\mu \phi(x) \rangle K^\mu \xrightarrow{\text{CPT}} -\langle \partial_\mu \phi(x) \rangle K^\mu$$

**So now:**


$$\mathcal{L}_{QED} \sim -\frac{1}{4} \mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu} + P_\mu A_\mu \tilde{\mathcal{F}}^{\mu\nu}$$

$P_\mu$ :  $P_0$  constant (explicit CPT violation)

$$\partial_\mu \phi(x) \longrightarrow \langle \partial_0 \phi(x) \rangle \text{ (cosmologically)}$$

$\partial_\mu f(R)$  (R: Ricci scalar)

# With the Chern-Simons term, the CMB polarization get rotated



$$Q^r \pm iU^r = e^{\pm 2i\bar{\alpha}}(Q \pm iU)$$

Rotation angle:  $\alpha = - \int_i^f dx^\mu p_\mu , \quad p_\mu \sim -\partial_\mu \phi$


$$C_\ell^{EE,r} = C_\ell^{EE} \cos^2(2\bar{\alpha}) + C_\ell^{BB} \sin^2(2\bar{\alpha})$$

$$C_\ell^{TE,r} = C_\ell^{TE} \cos(2\bar{\alpha})$$

$$C_\ell^{BB,r} = C_\ell^{EE} \sin^2(2\bar{\alpha}) + C_\ell^{BB} \cos^2(2\bar{\alpha})$$

$$C_\ell^{TB,r} = C_\ell^{TE} \sin(2\bar{\alpha})$$


$$C_\ell^{EB,r} = \frac{1}{2}(C_\ell^{EE} - C_\ell^{BB}) \sin(4\bar{\alpha}) ,$$



*Bo Feng, Mingzhe Li, Jun-Qing Xia,  
Xuelei Chen and Xinmin Zhang  
Phys. Rev. Lett. 96, 221302 (2006)*

# Measurements on $\bar{\alpha}$ from CMB experiments

Reference/Data	Rotation angle $\bar{\alpha}$
<sup>1</sup> WMAP3+BO3	$-6^\circ \pm 4^\circ$
<sup>2</sup> WMAP3	$-2.5^\circ \pm 3.0^\circ$
<sup>3</sup> WMAP5	$-1.7^\circ \pm 2.1^\circ$
<sup>4</sup> WMAP7	$-1.1^\circ \pm 1.4^\circ \pm 1.5^\circ$
<sup>5</sup> WMAP9	$-0.36^\circ \pm 1.24^\circ \pm 1.5^\circ$
<sup>6</sup> QUaD	$-0.56^\circ \pm 0.82^\circ \pm 0.5^\circ$
<sup>7</sup> BICEP1	$-2.60^\circ \pm 1.02^\circ$
<sup>8</sup> BICEP1	$-2.77^\circ \pm 0.86^\circ \pm 1.3^\circ$
<sup>9</sup> POLARBEAR	$-1.08^\circ \pm 0.20^\circ \pm 0.5^\circ$
<sup>10</sup> ACTPol	$-0.2^\circ \pm 0.5^\circ$
<sup>11</sup> Planck 2015	$0.35^\circ \pm 0.05^\circ \pm 0.28^\circ$



<sup>1</sup>Feng, et.al., 2006, PRL 96, 221302

<sup>2</sup>Cabella, et.al., J. 2007, PRD, 76, 123014

<sup>3</sup>Komatsu, et.al., 2009, ApJS, 180, 330-376

<sup>4</sup>Komatsu, et.al., 2011, ApJS, 192, 18

<sup>5</sup>Hinshaw, et.al., 2013, ApJS, 208, 19

<sup>6</sup>QUaD Collaboration, et.al., 2009, PRL, 102, 161302

<sup>7</sup>XIA, et.al., 2010, PLB 687, 129

<sup>8</sup>BICEP1 Collaboration, et.al., 2014, PRD, 89,062006

<sup>9</sup>The POLARBEAR Collaboration, et.al., 2014, ApJ, 794, 171

<sup>10</sup>Naess, et.al., 2014, JCAP, 10, 007

<sup>11</sup>Planck Collaboration, arXiv:1605.08633

**Cosmological *CPT* violating effect on CMB polarization**Mingzhe Li<sup>1,2,3</sup> and Xinmin Zhang<sup>4</sup>e-Print: [arXiv:0810.0403](https://arxiv.org/abs/0810.0403)<sup>1</sup>*Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany*<sup>2</sup>*Department of Physics, Nanjing University, Nanjing 210093, People's Republic of China*<sup>3</sup>*Joint Center for Particle, Nuclear Physics and Cosmology, Nanjing University–Purple Mountain Observatory, Nanjing 210093, People's Republic of China*<sup>4</sup>*Institute of High Energy Physics, Chinese Academy of Sciences, P.O. Box 918-4, Beijing 100049, People's Republic of China*  
(Received 15 October 2008; published 14 November 2008)

A dark energy scalar (or a function of the Ricci scalar) coupled with the derivative to the matter fields will violate the *CPT* symmetry during the expansion of the Universe. This type of cosmological *CPT* violation helps to generate the baryon number asymmetry and gives rise to the rotation of the photon polarization which can be measured in the astrophysical and cosmological observations, especially the experiments of the cosmic microwave background radiation. In this paper, we derive the rotation angle in a fully general relativistic way and present the rotation formulas used for the cosmic microwave background data analysis. Our formulas include the corrections from the spatial fluctuations of the scalar field. We also estimate the magnitude of these corrections in a class of dynamical dark energy models for quintessential baryo/leptogenesis.

DOI: [10.1103/PhysRevD.78.103516](https://doi.org/10.1103/PhysRevD.78.103516)

PACS numbers: 98.80.Cq, 98.70.Vc, 98.80.Es

$$C'^{EE}_l + C'^{BB}_l = \exp[-4C^\alpha(0)] \sum_{l'} \frac{2l' + 1}{4\pi} (C^{EE}_{l'} + C^{BB}_{l'}) \int_{-1}^1 d_{22}^{l'}(\beta) d_{22}^l(\beta) e^{4C^\alpha(\beta)} d\cos(\beta)$$

$$C'^{EE}_l - C'^{BB}_l = \cos(4\bar{\alpha}) \exp[-4C^\alpha(0)] \sum_{l'} \frac{2l' + 1}{4\pi} (C^{EE}_{l'} - C^{BB}_{l'}) \int_{-1}^1 d_{-22}^{l'}(\beta) d_{-22}^l(\beta) e^{-4C^\alpha(\beta)} d\cos(\beta)$$

$$C'^{EB}_l = \sin(4\bar{\alpha}) \exp[-4C^\alpha(0)] \sum_{l'} \frac{2l' + 1}{4\pi} (C^{EE}_{l'} - C^{BB}_{l'}) \int_{-1}^1 d_{-22}^{l'}(\beta) d_{22}^l(\beta) e^{-4C^\alpha(\beta)} d\cos(\beta)$$

$$C'^{TE}_l = C_l^{TE} \cos(2\bar{\alpha}) e^{-2C^\alpha(0)}$$

$$C'^{TB}_l = C_l^{TE} \sin(2\bar{\alpha}) e^{-2C^\alpha(0)}$$

*M. Li, B. Yu  
arXiv:1303.1881*

# How to Derotate the Cosmic Microwave Background Polarization

Marc Kamionkowski

e-Print: [arXiv:0810.1286](https://arxiv.org/abs/0810.1286) [*California Institute of Technology, Mail Code 130-33, Pasadena, California 91125, USA*

(Received 7 October 2008; revised manuscript received 12 February 2009; published 19 March 2009)

If the linear polarization of the cosmic microwave background is rotated in a frequency-independent manner as it propagates from the surface of last scatter, it may introduce a *B*-mode polarization. Here I show that measurement of higher-order *TE*, *EE*, *EB*, and *TB* correlations induced by this rotation can be used to reconstruct the rotation angle as a function of position on the sky. This technique can be used to distinguish primordial *B* modes from those induced by rotation. The effects of rotation can be distinguished geometrically from similar effects due to cosmic shear.

DOI: [10.1103/PhysRevLett.102.111302](https://doi.org/10.1103/PhysRevLett.102.111302)

PACS numbers: 98.70.Vc

## list of references on anisotropy of rotation angle:

- POLARBEAR Collaboration 1509.02461
- BICEP/Keck Collaboration 1705.02523
- Amit P.S. Yadav, Rahul Biswas, Meng Su, Matias Zaldarriaga 0902.4466
- Vera Gluscevic, Marc Kamionkowski, Asantha Cooray 0905.1687
- Vera Gluscevic, Duncan Hanson, Marc Kamionkowski, Christopher M. Hirata 1206.5546
- Carlo Cattaldi 1510.02629
- Toshiya Namikawa 1612.07855
- Dylan R. Sutton, Chang Feng, Christian L. Reichardt 1702.01871
- Dagoberto Contreras, Paula Boubel, Douglas Scott 1705.06387
- Levon Pogosian, Meir Shimon, Matthew Mewes, Brian Keating 1904.07855
- ...

# $(r, \bar{\alpha}, \delta\alpha)$ strongly correlated





FIG. 3.— Theoretical CMB power spectra (thick lines) for the best fit model obtained from the WMAP9+B03+BICEP2 data combination. The red thin line denotes the contribution of tensor perturbations with  $r = 0.2$ , and the black thin line is the total CMB BB power spectrum correspondingly. For comparison, we also show the BICEP2 observational data.

**Joint constraint on “B mode physics parameters” ( $r$ ,  $\bar{\alpha}$ ,  $\delta\alpha$ )  
with CMB polarization data  
from BICEP, Planck, ACT, SPT, POLARBEAR**



# AliCPT-1 collaboration

- Led by the IHEP, the international collaboration includes: about 100 scientists from China, US, Europe.
- Funded already by the CAS, NSF of China.
- Executive board: Xinmin Zhang (PI), Chao-Lin Kuo (US PI), Fang-jun Lu (Manager)



# **Core member institutes of AliCPT-1 collaboration**

- IHEP: Pipeline, data analysis, scan strategy, control system, site, mount, test/integration
- Stanford: Cryostat receiver, optics/AR, focal plane module
- NAOC: Logistics, site
- NIST: Detector arrays and modules, feedhorns and readout components
- ASU: LNAs, cryogenic harness, readout electronics
- NTU: Scan strategy, calibration
- APC: Science, data analysis
- USTC: E/B leakage, early universe models
- SJTU: Foregrounds, cross-correlations
- BNU: Foregrounds, lensing
- ...



# AliCPT-1 current status, future plan (AliCPT-2) and science


# AliCPT-1 Timeline & AliCPT-2

- 2014 – Project proposed
- 2017 – Site construction starts
- 2018 – Site construction ends
- 2019 – Cryostat under fab,  
design single pixel detector,  
mount under test,  
pipeline development
- 2019 – cryostat delivery at Stanford ,
- 2020 – Integration, test, commissioning at Stanford



**Deployment and first light with at least one and up to 4 modules by end of 2020**

## AliCPT-2:



- Add 15 modules to AliCPT-1 (with more frequencies?)  
2020 4 modules, 2021 10 modules, 2022 15 modules,  
2023 19 modules
- Large aperture Telescopes

# Mount



Sept. 2019


1. Bearing capacity
- max. 1.5tons



2. Driver spec. requirements
  - AZ: range  $\pm 270\text{deg}$ . angular speed  $>5\text{deg/sec}$
  - EL: range  $45\text{-}135\text{deg}$ . angular speed  $>1\text{deg/sec}$
  - DK: range  $0\text{-}181\text{deg}$ . angular speed  $>2\text{deg/sec}$
3. Pointing accuracy
  - better than  $1'$  w/o correction
  - Better than  $10''$  w/ pointing model correction



Test now, and expect to ship to B1-site soon.




Slides from Congzhan Liu

# AliCPT-1 in a nutshell


- 72cm aperture, FOV 20.8° (4-7det tiles)  
FOV 33.4° (19det tiles)
- 95/150GHz, 27/19% bandwidth
- 19' and 11'
- 1704 pol-sensitive, optical dichroich  
TESes per tile
- 280mK, NEP  $3\text{-}5 \sim 10^{-17}\text{W/sqrt(Hz)}$
- 4 detector modules
- Cryostat and optics: up to 19 det modules
- Forebaffle design: up to 7 det modules
- scanning in azimuth at constant elevation
- (45°-70°) elevation range
- up to 4°/s scanning speed
- Instrument design heritage BICEP3

Ahmed Z. et al., SPIE 2014



Slides from Maria Salatino

# Cryostat Receiver Overview



## Mechanics

- 300K, 50K, 4K tubes/baserings
- G-10CR structures

## Window

## IR Filters

- Zotefoam layers
- 50K Alumina/Nylon filters

## Cold Optics

- 4K Alumina lenses/AR epoxy mix

## Cryogenics

- 1/2x PT-420
- Custom GL10 absorption fridge

## TESes array and $\mu$ mux

14

Slides from Maria Salatino

# AliCPT sciences


## B mode physics:

- BB: Measuring the primordial gravitational waves (PGWs), tensor to scalar ratio ( $r$ ), probing the origin of the universe.
- TB, EB, BB: Measuring the rotation angle ( $\bar{\alpha}, \delta\alpha$ ), testing CPT symmetry.
- Investigating the CMB polarization hemispherical asymmetry.
- Studying the cross-correlation between AliCPT and DESI.
- Cosmology with precision measurement of CMB E modes.
- Studying the galactic foreground.
- Studying the dark energy property, neutrino masses


• • • • •

# Sensitivity

## Sensitivity on map level:




PLANCK 100 GHz & 143 GHz  
Pol ~50 uKarcmin in total




AliCPT 4 modules 1 season  
Median map-depth 14 uKarcmin


## Sensitivity on $r$ and $\bar{\alpha}$ :



according to PI's plan, AliCPT-1 for the first years' observation, AliCPT-1 + 2 for 2021-2023.



Raw sensitivity on  $r$



Raw sensitivity on  $\bar{\alpha}$

# Summary

AliCPT is one of the four gravitational waves projects in China (for the PGWs); The others being FAST, space probes TianQin & TaiJi (for the astrophysical GWs).

- AliCPT-1: First light in 2020.
- AliCPT-2: More detectors, more frequencies, larger aperture.
- Complimentary to SO in Atacama desert, BICEP Array in South Pole... for sky coverage, science (testing inflation theory, Chern-Simons theory, CPT symmetry...).

# Thanks!