


The background of the slide is a Cosmic Microwave Background (CMB) fluctuation map. It shows a complex pattern of temperature variations across the sky, with a color scale ranging from dark blue (cooler) to yellow and red (warmer). A prominent bright yellow/orange spot is visible in the upper right quadrant, and another smaller one is in the lower right. The overall texture is grainy and noisy, characteristic of CMB data.

Prompt cusps and the dark matter annihilation signal

Simon White
Max Planck Institute for Astrophysics



Prompt Cusps

- ...are relevant whenever $P(k)$ is sharply truncated at high k
- ...form promptly as each initial density peak collapses
- ...have density profiles, $\rho(r) \approx 24 \bar{\rho} (r / R)^{-1.5}$, where $\bar{\rho}$ is the mean cosmic DM density and $R = a_c(\delta / \nabla^2 \delta)^{1/2}$ is the size of the linear overdensity peak (both measured at t_c , the time of peak collapse)
- ...have, by $1.2 t_c$, mass, $M_{\text{cusp}} \sim 7 R^3 \bar{\rho}$, and size, $r_{\text{cusp}} \sim 0.1 R$
- ...have an inner core radius set by phase-space constraints, thus dependent on the cosmological origin of the DM
- ...suffer late-time tidal disruption only in star-dominated regions of galaxies (through encounters with individual stars)
- ...dominate the dark matter annihilation signal in all but the very densest regions of galaxies




BBKS-predicted peak and cusp distributions in Λ CDM

Delos+White 2023



$$m_{\chi} = 100 \text{ GeV}, \quad T_{\text{kd}} = 30 \text{ GeV}$$

$$J = \int \rho^2 dV$$




Growth with time
of the prompt cusp
population and its
annihilation signal

Milky Way annihilation radiation profiles


The profile due to cusps is much shallower than that due to the smoothly distributed dark matter

Cusp emission dominates at >1 kpc neglecting tides, at >3 kpc including the mean tide, and at >7 kpc including stellar encounters also


Prompt cusps do not affect the Fermi Galactic Centre Excess, but if this is due to annihilation then they contribute much of the $1 - 10$ GeV background



Stücker et al 2023



Bounds on the mass of a thermal WIMP



Curves are 95% upper bounds based on Fermi's measurement of the isotropic γ -ray background after subtraction of known source populations

Inclusion of prompt cusps raises the lower limit on mass by a factor of 150

Prompt cusps

- The origin and structure of prompt cusps differ from those of “normal” halos
- For a $m = 100$ GeV, $T_{\text{kd}} = 30$ GeV WIMP, prompt cusps have Earth mass and are a million times more abundant than Earth-mass planets in the Milky Way, accounting for a percent or two of all dark matter
- In the Milky Way they are significantly disrupted by tides and by stellar encounters at $r < 20$ kpc
- They have no observable dynamical or gravitational lensing effects
- They dominate the dark matter annihilation signal from the outer halo of the Milky Way and from all extragalactic objects, leading to a local luminosity density that is proportional to $\bar{\rho}_{\text{dm}}$ rather than $\bar{\rho}_{\text{dm}}^2$